

BALCH POND

2024 SAMPLING HIGHLIGHTS

Station Deep

Wakefield, NH

Acton and Newfield, ME



Water quality data displayed in Tables 1 and 2 are surface water measurements with the exception of the dissolved oxygen data that are collected near the lake bottom. Summary statistics are provided for samples collected between June 21 and September 16, 2024.

Blue = Oligotrophic

Light Green = Mesotrophic

Dark Green = Eutrophic

Gray = No Data




Figure 1. Balch Pond Water Quality (2024)

Table 1. 2024 Balch Pond Seasonal Averages and NH DES Aquatic Life Nutrient Criteria¹

Parameter	Oligotrophic "Excellent"	Mesotrophic "Fair"	Eutrophic "Poor"	Balch Pond Average (range)	Balch Pond Classification
Water Clarity (meters)	> 4.0 – 7.0	2.5 – 4.0	< 2.5	5.1 meters (3.7 – 6.2)	Oligotrophic
Chlorophyll <i>a</i> ¹ (ppb)	< 3.3	3.3 – 5.0	> 5.0 – 11.0	2.5 ppb (1.2 – 3.9)	Oligotrophic
Total Phosphorus ¹ (ppb)	< 8.0	8.0 – 12.0	> 12.0 – 28.0	7.4 ppb (5.2 – 8.5)	Oligotrophic
Dissolved Oxygen (ppm)	> 5.0 – 7.0	2.0 – 5.0	< 2.0	Not Assessed	Not Assessed

Table 2. 2024 Balch Pond Seasonal Average Accessory Water Quality Measurements

Parameter	Assessment Criteria					Balch Pond Average (range)	Balch Pond Classification
Color (color units)	< 10 uncolored	10 – 20 slightly colored	20 – 40 lightly tea colored	40 – 80 tea colored	> 80 highly colored	22.5 color units (range: 20.2 – 25.6)	Lightly tea colored
Alkalinity (ppm)	< 0.0 acidified	0.1 – 2.0 extremely vulnerable	2.1 – 10 moderately vulnerable	10.1 – 25.0 low vulnerability	> 25.0 not vulnerable	8.2 ppm (range: single value)	Moderately vulnerable
pH (std units)	< 5.5 suboptimal for successful growth and reproduction		6.5 – 9.0 optimal range for fish growth and reproduction			6.8 standard units (range: single value)	Optimal range for fish growth and reproduction
Specific Conductivity (uS/cm)	< 50 uS/cm Characteristic of minimally impacted NH lakes		50-100 uS/cm Lakes with some human influence	> 100 uS/cm Characteristic of lakes experiencing human disturbances		57.1 uS/cm (range: single value)	Characteristic of lakes with some human influence

Strategies to stabilize and improve water quality

Implement Best Management Practices (BMPs) within the Balch Pond watershed to minimize the adverse impacts of polluted runoff and erosion into Balch Pond. Refer to "[Landscaping at the Water's Edge: An Ecological Approach](#)" and "[New Hampshire Homeowner's Guide to Stormwater Management: Do-It-Yourself Stormwater Solutions for Your Home](#)" and the [Acton Wakefield Watershed Alliance webpages](#) for more information on how to reduce nutrient loading caused by overland runoff. NH Lakes also provides a series of resources aimed at educating residents and protecting our lakes and ponds.

Figure 2. Balch Pond (2024 Seasonal Data)
Secchi Disk Transparency and Chlorophyll *a* Data




Figure 3. Balch Pond (2024 Seasonal Data)
Secchi Disk Transparency and Dissolved Color Data




Figure 4. Balch Pond- Deep Site (1983-2024)
Long-term Secchi Disk Transparency and Chlorophyll *a* Data





Figure 5. Balch Pond - Deep Site (1983-2024)
Long-term Total Phosphorus and Dissolved Color Data



Figures 2 and 3. Seasonal comparison of Balch Pond water transparency (Secchi Disk depth), chlorophyll *a* and dissolved color for 2024. Shallower water transparency measurements oftentimes correspond to increases in chlorophyll *a* and/or color concentrations.

Figures 4 and 5. Annual median Balch Pond water transparency, chlorophyll *a*, dissolved color and total phosphorus concentrations measured between 1983 and 2024, through the New Hampshire Lakes Lay Monitoring Program and the New Hampshire Department of Environmental Services. The long-term data provide insight into the water quality fluctuations, among years, that have been documented in Balch Pond.

Figure 6. Balch Pond - Deep Site
Temperature Profiles (June 21, 2024)




Figure 7. Balch Pond - Deep Site
Dissolved Oxygen Profiles (June 21, 2024)




Figure 8. Balch Pond - 2022 Bi-weekly
Total Phosphorus Inter-site Comparison





Figure 9. Balch Pond - 2023 Bi-weekly
Total Phosphorus Inter-site Comparison




Figures 6 and 7. Balch Pond temperature and dissolved oxygen profiles collected on June 21, 2024.

Figures 8 and 9. Total phosphorus inter-comparison among sampling locations for the 2022 and 2023 seasons. All total phosphorus samples were collected in the warm surface water layer.

Figure 10. Balch Pond- 2024 Bi-weekly

Total Phosphorus Inter-site Comparison (June 21 through September 16, 2024)



Figures 10. Total phosphorus inter-comparison among sampling locations for the 2024 season. All total phosphorus samples were collected in the warm surface water layer.

Table 3. Saco River Watershed Lakes (Acton ME and Wakefield NH)
(2024 water quality data collected between May 13 and September 16)

Lake	Average (range) Secchi Disk Transparency (meters)	Average (range) Total Phosphorus (ppb)	Average (range) Chlorophyll- <i>a</i> (ppb)	Average (range) Dissolved Oxygen (ppm)
Balch Pond	5.1 meters (range: 3.7 – 6.2)	7.4 ppb (range: 5.2 – 8.5)	2.5 ppb (range: 1.2 – 3.9)	Not Assessed
Belleau Lake	1.8 meters (range: 1.4 – 2.2)	16.7 ppb (range: 16.1 – 17.3)	8.4 ppb (range: 5.9 – 12.9)	Not Assessed
Pine River Pond	6.3 meters (range: 5.0 – 7.2)	5.7 ppb (range: 5.2 – 6.7)	2.4 ppb (range: 1.9 – 2.7)	0.7 ppm (range: 0.0 – 2.9)
Province Lake	2.9 meters (range: 1.5 – 3.9)	17.5 ppb (range: 11.7 – 26.1)	4.8 ppb (range: 1.9 – 8.3)	Not Assessed

- Water quality data are reported for a deep reference sampling location in each water body
- Dissolved oxygen measurements were taken late season and from the bottom water layer
- “Not Assessed” indicates the site is too shallow to form a bottom water layer


Data Interpretation: Overview of factors to consider when reviewing the Balch Pond data

This highlight report provides a general overview of the current and historical conditions of Balch Pond. The report is intended to provide a simple assessment of the water quality trends. Should you have additional questions about interpreting your water quality results, we would be happy to discuss the data with you and/or any concerns you may have. In general, some factors that influence the current and long-term water quality results/trends for our New Hampshire lakes and ponds include:

- **Land-use Patterns** within the watershed (drainage basin) – Research indicates land use patterns have an impact on how much phosphorus (nutrient) is washing into our lakes. In general, more urbanized watersheds have a greater degree of phosphorus runoff than highly forested/vegetated drainage areas.
- **Weather Patterns** – Rainfall and temperature can influence water quality. Wet periods, and overland runoff, tend to be a time when elevated nutrients and other pollutants are transported into our lakes. Temperature can also influence water quality conditions since many aquatic plants and algae tend to respond to changing seasonal conditions. Unusually warm periods are sometimes tied to short-term algal and cyanobacteria blooms.
- **Best Management Practices (BMPs)** – The presence/absence of best management practices can have an interplay on water quality. BMPs are measures that are used to manage nutrients and other pollutants that could otherwise make their way into our lakes. Properties that employ BMPs, designed specifically to remove pollutants of concern (e.g. sediments and phosphorus), are less likely to contribute nutrients and other pollutants into our lakes.
- **Temperature (Thermal) Stratification** – Many lakes become thermally stratified during the summer months and may form three distinct thermal layers: upper water layer (epilimnion), middle lake layer (metalimnion) and bottom cold-water layer (hypolimnion). These thermal zones form a barrier to lake mixing, during the summer months, and can coincide with differences in dissolved oxygen and specific conductivity through the water column.
- **Internal Nutrient Loading** (nutrients that are introduced from the sediments along the lake bottom) – Some of our lakes experience significant internal nutrient loading. Such lakes generally tend to be well stratified and exhibit increasing deep water phosphorus concentrations, relative to surface levels.

Figure 11. Balch Pond

Acton & Newfield, ME and Wakefield, NH
2024 deep and shallow-water sampling locations



0 0.3 0.6 0.9 1.2 Miles

Aerial Orthophoto Source: Maine Office of GIS, National Agriculture Imagery Program 2015 Maine
GPS Coordinates collected by the UNH Center for Freshwater Biology



Extension

